Journal of MOLECULAR BIOLOGY

EDITOR-IN-CHIEF
J. C. KENDREW

ASSISTANT EDITOR
S. BRENNER

EDITORIAL BOARD
C. R. Cantor
P. Chambon
W. W. Franke
M. Gellert
G. A. Gilbert
M. Gottesman
R. Huber
H. E. Huxley
A. Klug
V. Luzzati
J. H. Miller
M. F. Moody
K. Simons

VOLUME 155

15 February to 15 March 1982

ACADEMIC PRESS
London New York San Francisco

Per
QH
301

Copyright © 1982, by Academic Press Inc. (London) Ltd.

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 155

Number 1, 15 February 1982

Kadesch, T. R., Rosenberg, S. and Chamberlin, M. J., Binding of *Escherichia coli* RNA Polymerase Holoenzyme to Bacteriophage T7 DNA. Measurements of Binding at Bacteriophage T7 Promoter A_1 Using a Template Competition Assay 1

Rosenberg, S., Kadesch, T. R. and Chamberlin, M. J., Binding of *Escherichia coli* RNA Polymerase Holoenzyme to Bacteriophage T7 DNA. Measurements of the Rate of Open Complex Formation at T7 Promoter A_1 31

Paul, C. H., Building Models of Globular Protein Molecules from their Amino Acid Sequences. I. Theory 53

Appendix: The Electrostatic Energy of Charges and Dipoles in a Spherical Dielectric Cavity 59

Zelwer, C., Risler, J. L. and Brunie, S., Crystal Structure of *Escherichia coli* Methionyl-tRNA Synthetase at 2·5 Å Resolution 63

Number 2, 25 February 1982

McCorkle, G. M. Altman, S., Large Deletion Mutants of *Escherichia coli* $tRNA_1^{Tyr}$ 83

Appendix: Nucleotide Sequences Flanking the $tRNA_1^{Tyr}$ Transcription Unit 98

Hirschel, B. J. and Berg, D. E., A Derivative of Tn*5* with Direct Terminal Repeats can Transpose 105

Oppenheim, A. B., Mahajna, G., Koby, S. and Altuvia, S., Regulation of the Establishment of Repressor Synthesis in Bacteriophage λ 121

Franks, N. P., Melchior, V., Kirschner, D. A. and Caspar, D. L. D., Structure of Myelin Lipid Bilayers. Changes During Maturation . . 133

Tauc, P., Leconte, C., Kerbiriou, D., Thiry, L. and Hervé, G., Coupling of Homotropic and Heterotropic Interactions in *Escherichia coli* Aspartate Transcarbamylase 155

LETTERS TO THE EDITOR

Kamiya, R. and Nagai, R., Structural Similarity Between Actin Bundles from Characean Algal Cells and Sea Urchin Oocytes 169

HENDERSON, D., GEISLER, N. and WEBER, K., A Periodic Ultrastructure in Intermediate Filaments 173

CIARROCCHI, G. and PEDRINI, A. M., Determination of Pyrimidine Dimer Unwinding Angle by Measurement of DNA Electrophoretic Mobility 177

NUMBER 3, 5 MARCH 1982

STÉVENIN, J., GATTONI, R., KEOHAVONG, P. and JACOB, M., Mild Nuclease Treatment as a Probe for a Non-random Distribution of Adenovirus-specific RNA Sequences and of Cellular RNA in Nuclear Ribonucleoprotein Fibrils 185

FRASER, N. W., BAKER, C. C., MOORE, M. A. and ZIFF, E. B., Poly(A) Sites of Adenovirus Serotype 2 Transcription Units 207

YANOFSKY, C. and VANCLEEMPUT, M., Nucleotide Sequence of *trpE* of *Salmonella typhimurium* and its Homology with the Corresponding Sequence of *Escherichia coli* 235

SCHIBLER, U., PITTET, A.-C., YOUNG, R. A., HAGENBÜCHLE, O., TOSI, M., GELLMAN, S. and WELLAUER, P. K., The Mouse α-Amylase Multigene Family. Sequence Organization of Members Expressed in the Pancreas, Salivary Gland and Liver 247

UDVARDY, A., SÜMEGI, J., TÓTH, É. C., GAUSZ, J., GYURKOVICS, H., SCHEDL, P. and ISH-HOROWICZ, D., Genomic Organization and Functional Analysis of a Deletion Variant of the 87A7 Heat Shock Locus of *Drosophila melanogaster* 267

ANDERSON, D. M., RICHTER, J. D., CHAMBERLIN, M. E., PRICE, D. H., BRITTEN, R. J., SMITH, L. D. and DAVIDSON, E. H., Sequence Organization of the Poly(A) RNA Synthesized and Accumulated in Lampbrush Chromosome Stage *Xenopus laevis* Oocytes. 281

WÜTHRICH, K., WIDER, G., WAGNER, G. and BRAUN, W., Sequential Resonance Assignments as a Basis for Determination of Spatial Protein Structures by High Resolution Proton Nuclear Magnetic Resonance 311

BILLETER, M., BRAUN, W. and WÜTHRICH, K., Sequential Resonance Assignments in Protein ^{1}H Nuclear Magnetic Resonance Spectra. Computation of Sterically Allowed Proton–Proton Distances and Statistical Analysis of Proton–Proton Distances in Single Crystal Protein Conformations 321

Appendix: Dependence of the short-range distances d_1, d_2 and d_3 on the dihedral angles ϕ_i, ψ_i and χ_i^1 341

WAGNER, G. and WÜTHRICH, K., Sequential Resonance Assignments in Protein ^{1}H Nuclear Magnetic Resonance Spectra. Basic Pancreatic Trypsin Inhibitor 347

WIDER, G., LEE, K. H. and WÜTHRICH, K., Sequential Resonance Assignments in Protein ^{1}H Nuclear Magnetic Resonance Spectra. Glucagon Bound to Perdeuterated Dodecylphosphocholine Micelles 367

LETTER TO THE EDITOR

ELLIOTT, G. F., SAYERS, Z. and TIMMINS, P. A., Neutron Diffraction Studies of the Corneal Stroma. 389

NUMBER 4, 15 MARCH 1982

CHAMPNESS, W. C. and SNYDER, L., The *gol* Site: A *Cis*-acting Bacteriophage T4 Regulatory Region that can affect Expression of all the T4 Late Genes 395

NAGASHIMA, H. and ASAKURA, S., Studies on Co-operative Properties of Tropomyosin–Actin and Tropomyosin–Troponin–Actin Complexes by the Use of *N*-Ethylmaleimide-treated and Untreated Species of Myosin Subfragment 1 409

Appendix: Application of Function γ 424

YUKIOKA, M., HATAYAMA, T. and INOUE, A., Two Species of Chromatin–RNA Polymerase II Complex Released from Rat Liver Nuclei by Nuclease Digestion 429

CLORE, G. M., GRONENBORN, A. M. and DAVIES, R. W., Theoretical Aspects of Specific and Non-specific Equilibrium Binding of Proteins to DNA as Studied by the Nitrocellulose Filter Binding Assay. Co-operative and Non-co-operative Binding to a One-dimensional Lattice 447

SQUIRE, J. M., HARFORD, J. J., EDMAN, A. C. and SJÖSTRÖM, M., Fine Structure of the A-band in Cryo-sections. III. Crossbridge Distribution and the Axial Structure of the Human C-zone 467

FERMI, G., PERUTZ, M. F., DICKINSON, L. C. and CHIEN, J. C. W., Structure of Human Deoxy Cobalt Haemoglobin 495

MORRIS, E. R., POWELL, D. A., GIDLEY, M. J. and REES, D. A., Conformation and Interactions of Pectins. I. Polymorphism between Gel and Solid States of Calcium Polygalacturonate 507

POWELL, D. A., MORRIS, E. R., GIDLEY, M. J. and REES, D. A., Conformations and Interactions of Pectins. II. Influence of Residue Sequence on Chain Association in Calcium Pectate Gels 517

LETTERS TO THE EDITOR

WILDE, C. D., CROWTHER, C. E. and COWAN, N. J., Isolation of a Multigene Family Containing Human α-Tubulin Sequences 533

MORNON, J. P., SURCOUF, E., BERTHOU, J., CORVOL, P. and FOOTE, S., Crystallization of Mouse Submaxillary Gland Renin 539

Digitized by the Internet Archive
in 2026

https://archive.org/details/journal-of-molecular-biology_february-15-march-15-1982_155_index-contents

Index to Volume 155

A-band fine structure in cryo-sections. Crossbridge distribution and the axial structure of the human C-zone, 467
Actin bundles from characean algal cells and sea urchin oocytes show structural similarity, 169
— complexes (see also Tropomyosin), 409
Adenovirus serotype type 2 transcription units, poly(A) sites, 207
— -specific RNA sequences and cellular RNA in nuclear ribonucleoprotein fibrils, mild nuclease treatment as a probe for a non-random distribution, 185
Algal cells and sea urchin oocytes, actin bundles show structural similarity, 169
ALTMAN, S. (see also MCCORKLE, G. M.), 83
ALTUVIA, S. (see also OPPENHEIM, A. B.), 121
Amino acid sequences, use in building models of globular protein molecules; theory, 53
α-Amylase multigene family of mouse. Sequence organization of members expressed in the pancreas, salivary gland and liver, 247
ANDERSON, D. M., RICHTER, J. D., CHAMBERLIN, M. E., PRICE, D. H., BRITTEN, R. J., SMITH, L. D. & DAVIDSON, E. H., Sequence organization of the poly(A) RNA synthesized and accumulated in lampbrush chromosome stage *Xenopus laevis* oocytes, 281
ASAKURA, S. (see also NAGASHIMA, H.), 409
Aspartate transcarbamylase, *Escherichia coli*, coupling of homotropic and heterotropic interactions, 155

Bacteriophage λ, regulation of the establishment of repressor synthesis, 121
— T4 regulatory region, *cis*-acting, that can affect expression of all the T4 late genes; the *gol* site, 395
— T7 DNA, binding of *Escherichia coli* RNA polymerase holoenzyme; measurements of binding at T7 promoter A_1 using a template competition assay, 1
———, ———————; ——— the rate of open complex formation at T7 promoter A_1, 31
BAKER, C. C. (see also FRASER, N. W.), 207
Basic pancreatic trypsin inhibitor; sequential resonance assignments in protein ^1H nuclear magnetic resonance spectra, 347
BERG, D. E. (see also HIRSCHEL, B. J.), 105
BERTHOU, J. (see also MORNON, J. P.), 539
BILLETER, M., BRAUN, W. & WÜTHRICH, K., Sequential resonance assignments in protein ^1H nuclear magnetic resonance spectra. Computation of sterically allowed proton–proton distances and statistical analysis of proton–proton distances in single crystal protein conformations, 321
Binding, equilibrium, of proteins to DNA as studied by the nitrocellulose filter binding assay, theoretical aspects; co-operative and non-co-operative binding to a one-dimensional lattice, 447
— of *Escherichia coli* RNA polymerase holoenzyme to bacteriophage T7 DNA. Measurements of binding at T7 promoter A_1 using a template competition assay, 1
———————. Measurements of the rate of open complex formation at T7 promoter A_1, 31
BRAUN, W. (see also BILLETER, M.), 321
—, — (see also WÜTHRICH, K.), 311
BRITTEN, R. J. (see also ANDERSON, D. M.), 281
BRUNIE, S. (see also ZELWER, C.), 63

Calcium pectate gels, influence of residue sequence on chain association; conformations and interactions of pectins, 517
— polygalacturonate, polymorphism between gel and solid states; conformations and interactions of pectins, 507
CASPAR, D. L. D. (see also FRANKS, N. P.), 133
Chain association in calcium pectate gels, influence of residue sequence; conformations and interactions of pectins, 517
CHAMBERLIN, M. E. (see also ANDERSON, D. M.), 281
—, M. J. (see also KADESCH, T. R.), 1
—, — (see also ROSENBERG, S.), 31

Champness, W. C. & Snyder, L., The *gol* site: a *cis*-acting bacteriophage T4 regulatory region that can affect expression of all the T4 late genes, 395
Characean algal cells and sea urchin oocytes, actin bundles show structural similarity, 169
Chien, J. C. W. (see also Fermi, G.), 495
Chromatin–RNA polymerase II complex, two species, released from rat liver nuclei by nuclease digestion, 429
Ciarrocchi, G. & Pedrini, A. M., Determination of pyrimidine dimer unwinding angle by measurement of DNA electrophoretic mobility, 177
cis-acting bacteriophage T4 regulatory region that can affect expression of all the T4 late genes; the *gol* site, 395
Clore, G. M., Gronenborn, A. M. & Davies, R. W., Theoretical aspects of specific and non-specific equilibrium binding of proteins to DNA as studied by the nitrocellulose filter binding assay. Co-operative and non-co-operative binding to a one-dimensional lattice, 447
Cobalt haemoglobin, deoxy, human: structure, 495
Conformations and interactions of pectins. Influence of residue sequence on chain association in calcium pectate gels, 517
— — — — —. Polymorphism between gel and solid states of calcium polygalacturonate, 507
Co-operative and non-co-operative binding to a one-dimensional lattice; theoretical aspects of equilibrium binding of proteins to DNA as studied by the nitrocellulose filter binding assay, 447
— properties of tropomyosin–actin and tropomyosin–troponin–actin complexes, studies by the use of *N*-ethylmaleimide-treated and untreated species of myosin subfragment 1, 409
Corneal stroma, neutron diffraction studies, 389
Corvol, P. (see also Mornon, J. P.), 539
Cowan, N. J. (see also Wilde, C. D.), 533
Crossbridge distribution and the axial structure of the human C-zone; fine structure of the A-band in cryo-sections, 467
Crowther, C. E. (see also Wilde, C. D.), 533
Cryo-sections, fine structure of the A-band; crossbridge distribution and the axial structure of the human C-zone, 467
Crystal structure of *Escherichia coli* methionyl-tRNA synthetase at 2·5 Å resolution, 63
Crystallization of mouse submaxillary gland renin, 539
C-zone axial structure and crossbridge distribution; fine structure of the A-band in cryo-sections, 467

Davidson, E. H. (see also Anderson, D. M.), 281
Davies, R. W. (see also Clore, G. M.), 447
Deletion mutants, large, of *Escherichia coli* $tRNA_1^{Tyr}$, 83
— variant of the 87A7 heat shock locus of *Drosophila melanogaster*, genomic organization and functional analysis, 267
Deoxy cobalt haemoglobin, human; structure, 495
DNA electrophoretic mobility, use in determination of pyrimidine dimer unwinding angle, 177
—, T7, binding of *Escherichia coli* RNA polymerase holoenzyme; measurements of binding at T7 promoter A_1, using a template competition assay, 1
—, —, — — — — — polymerase holoenzyme; measurements of the rate of open complex formation at T7 promoter A_1, 31
—, theoretical aspects of equilibrium binding of proteins as studied by the nitrocellulose filter binding assay; co-operative and non-co-operative binding to a one-dimensional lattice, 447
Dickinson, L. C. (see also Fermi, G.), 495
Drosophila melanogaster, genomic organization and functional analysis of a deletion variant of the 87A7 heat shock locus, 267

Edman, A. C. (see also Squire, J. M.), 467
87A7 heat shock locus of *Drosophila melanogaster*, genomic organization and functional analysis of a deletion variant, 267
Electrophoretic mobility of DNA, use in determination of pyrimidine dimer unwinding angle, 177
Electrostatic energy of charges and dipoles in a spherical dielectric cavity, 59
Elliott, G. F., Sayers, Z. & Timmins, P. A., Neutron diffraction studies of the corneal stroma, 389
Equilibrium binding of proteins to DNA as studied by the nitrocellulose filter binding assay, theoretical aspects; co-operative and non-co-operative binding to a one-dimensional lattice, 447
Escherichia coli aspartate transcarbamylase, coupling of homotropic and heterotropic interactions, 155
— — methionyl-tRNA synthetase, crystal structure at 2·5 Å resolution, 63

— — RNA polymerase holoenzyme, binding to bacteriophage T7 DNA. Measurements of binding at T7 promoter A_1 using a template competition assay, 1
— — — — —, — — — — —, — — the rate of open complex formation at T7 promoter A_1, 31
— — $tRNA_1^{Tyr}$ large deletion mutants, 83
— — *trpE* nucleotide sequence and its homology with the corresponding sequence of *Salmonella typhimurium*, 235
Expression of all the T4 late genes can be affected by a *cis*-acting bacteriophage T4 regulatory region; the *gol* site, 395

Fermi, G., Perutz, M. F., Dickinson, L. C. & Chien, J. C. W., Structure of human deoxy cobalt haemoglobin, 495
Filaments, intermediate, show a periodic ultrastructure, 173
Fine structure of the A-band in cryo-sections. Crossbridge distribution and the axial structure of the human C-zone, 467
Foote, S. (see also Mornon, J. P.), 539
Franks, N. P., Melchior, V., Kirschner, D. A. & Caspar, D. L. D., Structure of myelin lipid bilayers. Changes during maturation, 133
Fraser, N. W., Baker, C. C., Moore, M. A. & Ziff, E. B., Poly(A) sites of adenovirus serotype 2 transcription units, 207

Gattoni, R. (see also Stévenin, J.), 185
Gausz, J. (see also Udvardy, A.), 267
Geisler, N. (see also Henderson, D.), 173
Gellman, S. (see also Schibler, U.), 247
Gene expression, can be affected by a *cis*-acting bacteriophage T4 regulatory region; the *gol* site, 395
Genes, α-amylase, mouse: sequence organization expressed in the pancreas, salivary gland and liver, 247
Genomic organization and functional analysis of a deletion variant of the 87A7 heat shock locus of *Drosophila melanogaster*, 267
Gidley, M. J. (see also Morris, E. R.), 507
—, — (see also Powell, D. A.), 517
Globular proteins: building models of molecules from their amino acid sequences; theory, 53
Glucagon bound to perdeuterated dodecylphosphocholine micelles; sequential resonance assignments in protein ^1H nuclear magnetic resonance spectra, 367
gol site: a *cis*-acting bacteriophage T4 regulatory region that can affect expression of all the T4 late genes, 395
Gronenborn, A. M. (see also Clore, G. M.), 447
Gyurkovics, H. (see also Udvardy, A.), 267

Haemoglobin: structure of human deoxy cobalt haemoglobin, 495
Hagenbüchle, O. (see also Schibler, U.), 247
Harford, J. J. (see also Squire, J. M.), 467
Hatayama, T. (see also Yukioka, M.), 429
Heat shock locus 87A7 of *Drosophila melanogaster*, genomic organization and functional analysis of a deletion variant, 267
Henderson, D., Geisler, N. & Weber, K., A periodic ultrastructure in intermediate filaments, 173
Hervé, G. (see also Tauc, P.), 155
Heterotropic and homotropic interaction coupling in *Escherichia coli* aspartate transcarbamylase, 155
Hirschel, B. J. & Berg, D. E., A derivative of Tn*5* with direct terminal repeats can transpose, 105
Homotropic and heterotropic interaction coupling in *Escherichia coli* aspartate transcarbamylase, 155
Human deoxy cobalt haemoglobin structure, 495
— α-tubulin sequences contained in isolated multigene family, 533
— C-zone axial structure and crossbridge distribution; fine structure of the A-band in cryo-sections, 467

Inoue, A. (see also Yukioka, M.), 429
Ish-Horowicz, D. (see also Udvardy, A.), 267

Jacob, M. (see also Stévenin, J.), 185

KADESCH, T. R., ROSENBERG, S. & CHAMBERLIN, M. J., Binding of *Escherichia coli* RNA polymerase holoenzyme to bacteriophage T7 DNA. Measurements of binding at bacteriophage T7 promoter A_1 using a template competition assay, 1
—, — (see also ROSENBERG, S.), 31
KAMIYA, R. & NAGAI, R., Structural similarity between actin bundles from characean algal cells and sea urchin oocytes, 169
KEOHAVONG, P. (see also STÉVENIN, J.), 185
KERBIRIOU, D. (see also TAUC, P.), 155
KIRSCHNER, D. A. (see also FRANKS, N. P.), 133
KOBY, S. (see also OPPENHEIM, A. B.), 121

λ (see also Bacteriophage), 121
Lampbrush chromosome stage *Xenopus laevis* oocytes, sequence organization of the synthesized and accumulated poly(A) RNA, 281
LECONTE, C. (see also TAUC, P.), 155
LEE, K. H. (see also WIDER, G.), 367
Lipid bilayers of myelin, structure; changes during maturation, 133
Liver nuclei, rat, release two species of chromatin–RNA polymerase II complex by nuclease digestion, 429
—: sequence organization of the mouse α-amylase multigene family, 247

MCCORKLE, G. M., ALTMAN, S., Large deletion mutants of *Escherichia coli* $tRNA_1^{Tyr}$, 83
MAHAJNA, G. (see also OPPENHEIM, A. B.), 121
MELCHIOR, V. (see also FRANKS, N. P.), 133
Methionyl-tRNA synthetase from *Escherichia coli*, crystal structure at 2·5 Å resolution, 63
MOORE, M. A. (see also FRASER, N. W.), 207
MORNON, J. P., SURCOUF, E., BERTHOU, J., CORVOL, P. & FOOTE, S., Crystallization of mouse submaxillary gland renin, 539
MORRIS, E. R., POWELL, D. A., GIDLEY, M. J. & REES, D. A., Conformation and interactions of pectins. I. Polymorphism between gel and solid states of calcium polygalacturonate, 507
—, — (see also POWELL, D. A.), 517
Mouse α-amylase multigene family. Sequence organization of members expressed in the pancreas, salivary gland and liver, 247
— submaxillary gland renin crystallization, 539
Multigene family containing human α-tubulin sequences; isolation, 533
Mutants, deletion, large, of *Escherichia coli* $tRNA_1^{Tyr}$, 83
Myelin lipid bilayer structure; changes during maturation, 133
Myosin subfragment 1, *N*-ethylmaleimide-treated and untreated, use in studies on co-operative properties of tropomyosin–actin and tropomyosin–actin complexes, 409

NAGAI, R. (see also KAMIYA, R.), 169
NAGASHIMA, H. & ASAKURA, S., Studies on co-operative properties of tropomyosin–actin and tropomyosin–troponin–actin complexes by the use of *N*-ethylmaleimide-treated and untreated species of myosin subfragment 1, 409
N-ethylmaleimide-treated and untreated species of myosin subfragment 1, use in studies on co-operative properties of tropomyosin–actin and tropomyosin–actin complexes, 409
Neutron diffraction studies of the corneal stroma, 389
Nitrocellulose filter binding assay studies of equilibrium binding of proteins to DNA, theoretical aspects; co-operative and non-co-operative binding to a one-dimensional lattice, 447
Nuclear ribonucleoprotein fibrils, mild nuclease treatment as a probe for a non-random distribution of adenovirus-specific RNA sequences and of cellular RNA, 185
Nuclease digestion releases two species of chromatin–RNA polymerase II complex from rat liver nuclei, 429
— treatment, mild, as a probe for a non-random distribution of adenovirus-specific RNA sequences and of cellular RNA in nuclear ribonucleoprotein fibrils, 185

OPPENHEIM, A. B., MAHAJNA, G., KOBY, S. & ALTUVIA, S., Regulation of the establishment of repressor synthesis in bacteriophage λ, 121

Pancreas: sequence organization of the mouse α-amylase multigene family, 247
PAUL, C. H., Building models of globular protein molecules from their amino acid sequences. I. Theory, 53
Pectins: conformations and interactions; influence of residue sequence on chain association in calcium pectate gels, 517
—: — — —; polymorphism between gel and solid states of calcium polygalacturonate, 507
PEDRINI, A. M. (see also CIARROCCHI, G.), 177
Periodic ultrastructure in intermediate filaments, 173
PERUTZ, M. F. (see also FERMI, G.), 495
PITTET, A.-C. (see also SCHIBLER, U.), 247
Poly(A) RNA synthesized and accumulated in lampbrush chromosome stage *Xenopus laevis* oocytes, sequence organization, 281
— sites of adenovirus serotype 2 transcription units, 207
POWELL, D. A., MORRIS, E. R., GIDLEY, M. J. & REES, D. A., Conformations and interactions of pectins. II. Influence of residue sequence on chain association in calcium pectate gels, 517
—, — (see also MORRIS, E. R.), 507
PRICE, D. H. (see also ANDERSON, D. M.), 281
Promoter A_1 on bacteriophage T7 DNA, measurements of binding of *Escherichia coli* RNA polymerase holoenzyme using a template competition assay, 1
— — — — — —, — — the rate of open complex formation with *Escherichia coli* RNA polymerase holoenzyme, 31
Proton nuclear magnetic resonance, sequential resonance assignments as a basis for a determination of spatial protein structures, 311
— — — —, — — —; basic pancreatic trypsin inhibitor, 347
— — — —, — — —; computation of sterically allowed proton–proton distances and statistical analysis of proton–proton-distances in single crystal protein conformations, 321
— — — —, — — —; glucagon bound to perdeuterated dodecylphoscholine micelles, 367
Proton–proton distances in protein conformations, 321
Protein equilibrium binding to DNA as studied by the nitrocellulose filter binding assay, theoretical aspects; co-operative and non-co-operative binding to a one-dimensional lattice, 447
— conformations by high resolution proton nuclear magnetic resonance, sequential resonance assignments as a basis for determination, 311
— proton nuclear magnetic resonance spectra, sequential resonance assignments; basic pancreatic trypsin inhibitor, 347
— — — — — —, — — —; computation of sterically allowed proton–proton distances and statistical analysis of proton–proton distances in single crystal protein conformations, 321
— — — — — —, — — —; glucagon bound to perdeuterated dodecylphosphocholine micelles, 367
Pyrimidine dimer unwinding angle, determination by measurement of DNA electrophoretic mobility, 177

Rat liver nuclei release two species of chromatin–RNA polymerase II complex by nuclease digestion, 429
REES, D. A. (see also MORRIS, E. R.), 507
—, — (see also POWELL, D. A.), 517
Regulation of the establishment of repressor synthesis in bacteriophage λ, 121
Renin crystallization, after isolation from mouse submaxillary gland, 539
Repressor synthesis in bacteriophage λ, regulation of its establishment, 121
RNA, poly(A), synthesized and accumulated in lampbrush chromosome stage *Xenopus laevis* oocytes, sequence organization, 281
— polymerase complex (see also Chromatin), 429
— — holoenzyme, *Escherichia coli*, binding to bacteriophage T7 DNA; measurements of binding at T7 promoter A_1, using a template competition assay, 1
— — —, — —, — — — — —; measurements of the rate of open complex formation at T7 promoter A_1, 31
— sequences, adenovirus-specific, and cellular RNA in nuclear ribonucleoprotein fibrils, mild nuclease treatment as a probe for a non-random distribution, 185
Ribonucleoprotein fibrils, mild nuclease treatment as a probe for a non-random distribution of adenovirus-specific RNA sequences and of cellular RNA, 185
RICHTER, J. D. (see also ANDERSON, D. M.), 281
RISLER, J. L. (see also ZELWER, C.), 63

ROSENBERG, S., KADESCH, T. R. & CHAMBERLIN, M. J., Binding of *Escherichia coli* RNA polymerase holoenzyme to bacteriophage T7 DNA. Measurements of the rate of open complex formation at T7 promoter A_1, 31
—, — (see also KADESCH, T. R.), 1

Salivary gland: sequence organization of the mouse α-amylase multigene family, 247
Salmonella typhimurium trpE nucleotide sequence and its homology with the corresponding sequence of *Escherichia coli*, 235
SAYERS, Z. (see also ELLIOTT, G. F.), 389
SCHEDL, P. (see also UDVARDY, A.), 267
SCHIBLER, U., PITTET, A.-C., YOUNG, R. A., HAGENBÜCHLE, O., TOSI, M., GELLMAN, S. & WELLAUER, P. K., The mouse α-amylase multigene family. Sequence organization of members expressed in the pancreas, salivary gland and liver, 247
Sea urchin oocytes and characean algal cells, actin bundles show structural similarity, 169
Sequence, nucleotide, of *trpE* of *Salmonella typhimurium* and its homology with the corresponding sequence of *Escherichia coli*, 235
— organization of mouse α-amylase multigene family expressed in the pancreas, salivary gland and liver, 247
— organization of the poly(A) RNA synthesized and accumulated in lampbrush chromosome stage *Xenopus laevis* oocytes, 281
Sequences, amino acid, use in building models of globular protein molecules; theory, 53
—, nucleotide, flanking the $tRNA_1^{Tyr}$ gene transcription unit, 98
—, RNA, adenovirus-specific, and cellular RNA in nuclear ribonucleoprotein fibrils, mild nuclease treatment as a probe for a non-random distribution, 185
SJÖSTRÖM, M. (see also SQUIRE, J. M.), 467
SMITH, L. D. (see also ANDERSON, D. M.), 281
SNYDER, L. (see also CHAMPNESS, W. C.), 395
SQUIRE, J. M., HARFORD, J. J., EDMAN, A. C. & SJÖSTRÖM, M., Fine structure of the A-band in cryosections. III. Crossbridge distribution and the axial structure of the human C-zone, 467
STÉVENIN, J., GATTONI, R., KEOHAVONG, P. & JACOB, M., Mild nuclease treatment as a probe for a non-random distribution of adenovirus-specific RNA sequences and of cellular RNA in nuclear ribonucleoprotein fibrils, 185
Structural similarity between actin bundles from characean algal cells and sea urchin oocytes, 169
Structure of *Escherichia coli* methionyl-tRNA synthetase at 2·5 Å resolution, 63
— — human deoxy cobalt haemoglobin, 495
— — myelin lipid bilayers; changes during maturation, 133
Structures of protein by high resolution proton nuclear magnetic resonance, sequential resonance assignments as a basis for determination, 311
— — — — — — — — — —, — — —; computation of sterically allowed proton–proton distances and statistical analysis of proton–proton distances in single crystal protein conformations, 321
Submaxillary gland renin of mouse; crystallization, 539
SÜMEGI, J. (see also UDVARDY, A.), 267
SURCOUF, E. (see also MORNON, J. P.), 539

T4 (see also Bacteriophage), 395
T7 (see also Bacteriophage), 1, 31
TAUC, P., LECONTE, C., KERBIRIOU, D., THIRY, L. & HERVÉ, G., Coupling of homotropic and heterotropic interactions in *Escherichia coli* aspartate transcarbamylase, 155
Template competition assay, use in measurements of binding of *Escherichia coli* RNA polymerase holoenzyme at bacteriophage T7 promoter A_1, 1
THIRY, L. (see also TAUC, P.), 155
TIMMINS, P. A. (see also ELLIOTT, G. F.), 389
Tn*5* derivative with direct terminal repeats can transpose, 105
TOSI, M. (see also SCHIBLER, U.), 247
TÓTH, É. C. (see also UDVARDY, A.), 267
Transfer RNA_1^{Tyr}, *Escherichia coli*, large deletion mutants, 83
— — gene transcription unit, flanking nucleotide sequences, 98
Transcription units, adenovirus serotype 2, poly(A) sites, 207
— by a derivative of Tn*5* with direct terminal repeats, 105

Tropomyosin–actin and tropomyosin–troponin–actin complexes, studies on co-operative properties using *N*-ethylmaleimide-treated and untreated species of myosin subfragment 1, 409
Troponin complexes (see also Tropomyosin), 409
trpE nucleotide sequence of *Salmonella typhimurium* and its homology with the corresponding sequence of *Escherichia coli*, 235
α-Tubulin sequences, human, contained in isolated multigene family, 533

Udvardy, A., Sümegi, J., Tóth, É. C., Gausz, J., Gyurkovics, H., Schedl, P. & Ish-Horowicz, D., Genomic organization and functional analysis of a deletion variant of the 87A7 heat shock locus of *Drosophila melanogaster*, 267
Ultrastructure, periodic, in intermediate filaments, 173
Unwinding angle, pyrimidine dimer, determination by measurement of DNA electrophoretic mobility, 177

Vancleemput, M. (see also Yanofsky, C.), 235

Wagner, G. & Wüthrich, K., Sequential resonance assignments in protein ^{1}H nuclear magnetic resonance spectra. Basic pancreatic trypsin inhibitor, 347
—, — (see also Wüthrich, K.), 311
Weber, K. (see also Henderson, D.), 173
Wellauer, P. K. (see also Schibler, U.), 247
Wider, G., Lee, K. H. & Wüthrich, K., Sequential resonance assignments in protein ^{1}H nuclear magnetic resonance spectra. Glucagon bound to perdeuterated dodecylphosphocholine micelles, 367
—, — (see also Wüthrich, K.), 311
Wilde, C. D., Crowther, C. E. & Cowan, N. J., Isolation of a multigene family containing human α-tubulin sequences, 533
Wüthrich, K., Wider, G., Wagner, G. & Braun, W., Sequential resonance assignments as a basis for determination of spatial protein structures by high resolution proton nuclear magnetic resonance, 311
—, — (see also Billeter, M.), 321
—, — (see also Wagner, G.), 347
—, — (see also Wider, G.), 367

Xenopus laevis: sequence organization of the poly(A) RNA synthesized and accumulated in lampbrush chromosome stage oocytes, 281

Yanofsky, C. & Vancleemput, M., Nucleotide sequence of *trpE* of *Salmonella typhimurium* and its homology with the corresponding sequence of *Escherichia coli*, 235
Young, R. A. (see also Schibler, U.), 247
Yukioka, M., Hatayama, T. & Inoue, A., Two species of chromatin–RNA polymerase II complex released from rat liver nuclei by nuclease digestion, 429

Zelwer, C., Risler, J. L. & Brunie, S., Crystal structure of *Escherichia coli* methionyl-tRNA synthetase at 2·5 Å resolution, 63
Ziff, E. B. (see also Fraser, N. W.), 207